


6 to 12 Months



6 TO 12 MONTHS

Safety for Your Child

Did you know that hundreds of children younger than 1 year die every year in the United States because of injuries — most of which can be prevented?

Often, injuries happen because parents are not aware of what their children can do. Your child is a fast learner and will suddenly be able to *roll over*, *crawl*, *sit*, and *stand*. Your child may *climb* before walking, or *walk* with support months before you expect. Your child will *grasp* at almost anything and reach things they could not reach before.

Falls

Because of your child's new abilities, he or she will fall often. Protect your child from injury. **Use gates on stairways and doors. Install operable window guards** on all windows above the first floor. **Remove sharp-edged or hard furniture** from the room where your child plays.

Do not use a baby walker. Your child may tip it over, fall out of it, or fall down the stairs in it. Baby walkers allow children to get to places where they can pull hot foods or heavy objects down on themselves.

If your child has a serious fall or does not act normally after a fall, call your doctor.

Burns

At 6 to 12 months children grab at everything. **NEVER** leave cups of hot coffee on tables or counter edges. **And NEVER carry hot liquids or food near your child or while holding your child.** He or she could get burned. Also, if your child is left to crawl or walk around stoves, wall or floor heaters, or other hot appliances, he or she is likely to get burned. **A safer place for your child** while you are cooking, eating, or unable to provide your full attention is the **playpen, high chair, or crib.**

If your child does get burned, put cold water on the burned area immediately. Then cover the burn loosely with a bandage or clean cloth. Call your doctor for all burns. To protect your child from tap water scalds, reduce the temperature of your hot water heater to a maximum of 120°F.

Test the batteries on your smoke alarm every month to be sure that they work. Change the batteries at least once a year on a date you'll remember, such as daylight savings time.



(over)

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



SUPPORTED BY A GRANT FROM PFIZER CONSUMER HEALTHCARE, MAKERS OF NEOSPORIN®

Drowning

At this age your child loves to play in water. Empty all the water from a bathtub, pail, or any container of water immediately after use. Keep the door to the bathroom closed. **NEVER leave your child alone in or near a bathtub, pail of water, wading or swimming pool, or any other water, even for a moment.** Drowning can happen in less than 2 inches of water. Knowing how to swim does NOT make your child water safe at this age. Stay within an arm's length of your child around water.

If you have a swimming pool, now is the time to **install a fence** that separates the house from the pool. The pool should be fenced in on all 4 sides. Most children drown because they fall into a pool that is not fenced off from the house. Be prepared — install a fence around your pool now, before your child begins to walk!



Poisoning and Choking

Your child will explore the world by *putting anything and everything into his or her mouth*. NEVER leave small objects or balloons in your child's reach, even for a moment. Don't feed your child hard pieces of food such as hot dogs, raw carrots, grapes, peanuts, or popcorn. Cut all his or her food into thin slices to prevent choking.

Be prepared if your child starts to choke. Learn how to save the life of a choking child. Ask your doctor to recommend the steps you need to take.

Children will put everything into their mouths, even if it doesn't taste good. Many ordinary things in your house **can be poisonous** to your child. Be sure to keep household products such as cleaners, chemicals, and medicines up, up, and away, completely out of sight and reach. Never store lye drain cleaners in your home. **Use safety latches** on drawers and cupboards. Remember, your child doesn't understand or remember "no" while exploring.

If your child does eat something that could be poisonous, call your doctor or the Poison Center immediately. Have syrup of ipecac on hand to make your child vomit, but use it only if you are told to do so by the Poison Center or your doctor.



Strangulation and Suffocation

Place your baby's crib away from windows. **Cords from window blinds and draperies can strangle your child.** Tie cords high and out of reach. Do not knot cords together.

Plastic wrappers and bags form a tight seal if placed over the mouth and nose and may suffocate your child. Keep them away from your child.

And Remember Car Safety

Car crashes are still a **great danger** to your child's life and health. Most injuries and deaths caused by car crashes **can be prevented** by the use of car safety seats EVERY TIME your child is in the car. An infant must always ride in a rear-facing car safety seat in the back seat until he or she is at least 1 year of age and at least 20 pounds. A rear-facing car safety seat should NEVER be placed in front of a passenger air bag. Your child, besides being much safer in a car safety seat, will behave better so you can pay attention to your driving. **The safest place for all infants and children to ride is in the back seat.**

Do not leave your child alone in a car. Keep vehicles and their trunks locked. Death from excess heat may occur in a closed car in warm weather in a short time.

Remember, the biggest threat to your child's life and health is an injury.

