

Well Child Care and Immunization Schedule

Our clinic follows the preventive care schedule recommended by the American Academy of Pediatrics and is accepted by all major health insurance plans. Each visit will include a complete physical exam, review of developmental milestones, diet, safety precautions, and behavioral management for the respective ages.

- Birth – Your baby will be seen daily by the pediatrician with the first visit usually within 12 hours of birth. If a circumcision is desired it is usually performed on the second day. Hepatitis B (HBV) #1 is given in the hospital.
- 3-7 day Physical, bilirubin lab if jaundice is significant
- 2 week Physical, neonatal screen
- 1 Month – Physical, HBV #2, neonatal screen if not completed
- 2 Months – Physical, Diphtheria / Tetanus / acellular Pertussis / Haemophilus Influenza B / inactivated Polio (Pentacel) #1, Streptococcus pneumoniae-Prevnar (SPV) #1, Rotavirus (RV) #1
- 4 Months – Physical, Pentacel #2, SPV #2, RV #2
- 6 Months – Physical, Pentacel #3, SPV #3, Flu in season
- 9 Months – Physical, HBV #3, complete blood count, lead count , Flu in season
- 1 Year – Physical, Measles / Mumps / Rubella (MMR) #1, Varicella-chicken pox, SPV #4, Flu in season, infant/toddler developmental screen
- 15 Months – Physical, DTaP #4, Hib #4 Hepatitis A #1 (HAV), flu in season, blood count, lead count
- 18 Months – Physical , CHAT Autism screen , flu in season
- 2 Years – Physical, Hepatitis A #2, blood count, lead count , vision screen ,flu in season
- 3 Years – Physical, vision screen, blood count, lead count , flu in season
- 4 Years – Physical, DTaP #5, IPV #4, MMR#2, Varicella-chicken pox #2, vision test, hearing test , flu in season, blood count, lead count
- 5 Years – Physical, vision test, hearing test, (if not done at 4 year), flu in season
- 6 - 24 Years -Physical each year, flu in season, complete blood count and lipid panel every 2 years additional labs as needed
- 11 Years - diphtheria-tetanus-pertussis booster (Tdap), Meningococcal vaccine, Human Papillomavirus (Gardasil/Cerarix) 3 doses
- 16-18 Years Meningococcal vaccine #2

Flu Vaccine is recommended for everyone 6 months and older, not just high risk patients.

Updated vaccine information is under the immunization tab, the CDC Vaccine Information Sheets are there for review.

PEDS Developmental Screen 2 - 8 years

