

Children's **TYLENOL**[®]

- The #1 branded fever reducer recommended by Pediatricians¹
- The fever reducer medicine with 0-6 month professional dosing



		Children's Tylenol[®] Infants' Drops 80 mg/0.8 mL	Children's Tylenol[®] Suspension Liquid 160 mg/5 mL	Children's Tylenol[®] Soft Chews Chewable Tablets 80 mg each	Junior Strength Tylenol[®] Chewable Tablets 160 mg each
Dose		mL	Teaspoon (tsp)*	Tablet	Tablet
WEIGHT	AGE	(Use only the dropper provided)	(Use only the dosing cup provided)		
6-11 lbs	0-3 mos	0.4mL			
12-17 lbs	4-11 mos	0.8mL	1/2 (tsp)		
18-23 lbs	12-23 mos	1.2mL (0.8 + 0.4mL)	3/4 (tsp)		
24-35 lbs	2-3 yrs	1.6mL (0.8 + 0.8mL)	1 (tsp)	2	
36-47 lbs	4-5 yrs		1-1/2 (tsp)	3	
48-59 lbs	6-8 yrs		2 (tsp)	4	2
60-71 lbs	9-10 yrs		2 1/2 (tsp)	5	2 1/2
72-95 lbs	11 yrs		3 (tsp)	6	3
96 lbs & over	12 yrs				4

Use only as directed.

NOTE: If possible, use weight to dose; otherwise use age. To arrive at the correct dose, weigh your child before giving TYLENOL[®]. All dosages may be repeated every 4 hours, but not more than 5 times daily.

A healthcare professional should be consulted for dosing for children under the age of two years.

WARNINGS:

- Children's TYLENOL[®] should not be taken for pain for more than 5 days or for fever for more than 3 days unless directed by a physician. If pain or fever persists or gets worse, if new symptoms occur, or if redness or swelling is present, a physician should be consulted because these could be signs of a serious condition.
- Do not exceed recommended dose. Taking more than the recommended dose (even once) may not provide more relief and could cause serious health problems. Keep this and all drugs out of the reach of children. In case of accidental overdose, contact a physician or poison control center immediately. Prompt medical attention is critical even if you do not notice any signs or symptoms.
- Do not use with any other product containing acetaminophen.
- Do not use Adult Extra Strength TYLENOL[®] products for children under 12 years of age.
- Do not use a kitchen teaspoon. Only use the dosing cup provided.

Children's **Motrin**[®]

- Pediatricians' #1 choice for fever relief that lasts up to 8 hours¹
- The #1 ibuprofen choice of consumers by more than 3 to 1 over Children's Advil[®] 2



		Children's Motrin[®] Concentrated Infants' Drops 50 mg/1.25 mL	Children's Motrin[®] Suspension 100 mg/5 mL	Children's Motrin[®] Chewable Tablets 50 mg	Junior Strength Motrin[®] Chewable Tablets 100 mg	Junior Strength Motrin[®] Caplets 100 mg
Dose		mL	Teaspoon (tsp)*	Tablet	Tablet	Caplet
WEIGHT	AGE	(Use only the dropper provided)	(Use only the dosing cup provided)			
Under 6 mos		Consult Your Child's Doctor				
12-17 lbs	6-11 mos	1.25 mL	---	---	---	---
18-23 lbs	12-23 mos	1.875 mL	---	---	---	---
24-35 lbs	2-3 yrs	---	1 (tsp)	2 tablets	---	---
36-47 lbs	4-5 yrs	---	1-1/2 (tsp)	3 tablets	---	---
48-59 lbs	6-8 yrs	---	2 (tsp)	4 tablets	2 tablets	2 caplets
60-71 lbs	9-10 yrs	---	2 1/2 (tsp)	5 tablets	2 1/2 tablets	2 1/2 caplets
72-95 lbs	11 yrs	---	3 (tsp)	6 tablets	3 tablets	3 caplets
One Dose Lasts 6-8 Hours						

Children's Motrin is available over-the-counter (OTC) for the temporary reduction of fever and relief of minor aches and pains due to colds, flu, sore throat, headaches and toothaches.

OTC Dosing: The recommended dose is 7.5 mg/kg every 6-8 hours. The recommended maximum daily dose is 30 mg/kg.

Keep all medication out of the reach of children. Use only as directed.

*Do not use a kitchen teaspoon. Only use the dosing cup provided.

References: 1. IMS NDI Pediatricians' recommendations, 12 months ending August 2002.
2. IRI Data, 52 weeks ending September 22, 2002.



McNEIL CONSUMER & SPECIALTY PHARMACEUTICALS
DIVISION OF McNEIL-PPC, INC.
FORT WASHINGTON, PA 19034 USA

OTCM-0032-P

Most widely treated ailments



- Everyday fevers
- Teething pain
- Immunizations
- Headaches
- Minor aches and pains



- High fevers
- Headaches
- Sprains
- Strains

Tips on Giving Medicine Correctly...



- **Always use the dosing device provided with the product to give your child medicine.** Household items such as measuring spoons are a less accurate way to ensure that you're giving the proper dose.
- **Make sure that all caregivers are aware of the medication you've given your child.** Avoid extra dosing by a parent or sitter who's unaware of what your child's already taken.
- **Never give adult medicine to a child.** Though it may seem that you can calculate a proper dosage of an adult medicine based on your child's size, medicines come in different concentrations designed to best treat infants, children and adults.
- **Check labels to be sure that different medicines aren't combining to provide too high a dose.** Cold medicines and pain relievers often have some of the same ingredients; a full dose of each may be twice as much medicine as is recommended.
- **Keep a log of your child's daily medications.** Have a pad and paper handy to write down exactly when and how much medicine is given throughout the day to ensure proper dosing.

If you have questions or comments, visit our website at www.GetWellKids.com or call your Healthcare Professional